arXiv:1610.07042v3 [math.GR] 20 Nov 2016

# FINITE $p$-GROUPS HAVING SCHUR MULTIPLIER OF MAXIMUM ORDER

SUMANA HATUI

ABSTRACT. Let $G$ be a non-abelian $p$-group of order $p^n$ and $M(G)$ denote the Schur multiplier of $G$. Niroomand proved that $|M(G)| \leq p^{\frac{1}{2}(n+k-2)(n-k-1)+1}$ for non-abelian $p$-groups $G$ of order $p^n$ with derived subgroup of order $p^k$. Recently Rai classified $p$-groups $G$ of nilpotency class 2 for which $|M(G)|$ attains this bound. In this article we show that there is no finite $p$-group $G$ of nilpotency class $c \geq 3$ for $p \neq 3$ such that $|M(G)|$ attains this bound. Hence $|M(G)| \leq p^{\frac{1}{2}(n+k-2)(n-k-1)}$ for $p$-groups $G$ of class $c \geq 3$ where $p \neq 3$. We also construct a $p$-group $G$ for $p = 3$ such that $|M(G)|$ attains the Niroomand's bound.

## 1. INTRODUCTION

Let $G$ be a finite $p$-group and $G'$ denotes its derived subgroup. The concept of Schur multiplier $M(G)$ of a group $G$ was introduced by Schur [1] in 1904 while studying of projective representation of groups. In 1956, Green [2] proved that $|M(G)| \leq p^{\frac{1}{2}n(n-1)}$ for $p$-groups $G$ of order $p^n$. So $|M(G)| = p^{\frac{1}{2}n(n-1)-t(G)}$, for some $t(G) \geq 0$. It is of interest to characterize the structure of all non-abelian $p$-groups $G$ by the order of the Schur multiplier $M(G)$, i.e., when $t(G)$ is known. This problem was studied by several authors for various values of $t(G)$ and the results are known for $0 \leq t(G) \leq 6$ in [3, 4, 5, 7, 8, 12].

Later Niroomand [6] improved the Green's bound and showed that for non abelian $p$-groups $G$ of order $p^n$ with $|G'| = p^k$,

$$|M(G)| \leq p^{\frac{1}{2}(n+k-2)(n-k-1)+1}.$$

He also classified groups $G$ such that $|M(G)| = p^{\frac{1}{2}(n+k-2)(n-k-1)+1}$ with $k = 1$. For our convention instead of writing $|M(G)| = p^{\frac{1}{2}(n+k-2)(n-k-1)+1}$ we shall write $|M(G)|$ attains the bound, throughout this paper.

Recently Rai [16, Theorem 2.1] classified finite $p$-groups $G$ of class 2 such that $|M(G)|$ attains the bound.

Aim of this paper is to continue this line of investigation and to look into the classification of arbitrary finite $p$-groups attending this bound. It, surprisingly turns out that for $p \neq 3$ there is no finite $p$-group $G$ of nilpotency class $c \geq 3$ such that $|M(G)|$ attains the bound. Hence for $p$-groups $G$ of class $\geq 3$ and $p \neq 3$ we improve

2010 *Mathematics Subject Classification.* 20J99, 20D15.
*Key words and phrases.* Schur Multiplier, Finite $p$-group.

the bound and in this case $|M(G)| \leq p^{\frac{1}{2}(n+k-2)(n-k-1)}$ where $|G'| = p^k$. Now one can ask what will happen for $p = 3$? Is this above statement true for $p = 3$?
The answer to this question is no. We construct an example, see Lemma 3.4, which gives the answer of this question. Now we state our main result which is the following.

**Theorem 1.1.** *There is no non-abelian $p$-group $G$ of order $p^n$, $p \neq 3$, having nilpotency class $c \geq 3$ with $|G'| = p^k$ and $|M(G)| = p^{\frac{1}{2}(n+k-2)(n-k-1)+1}$. In particular, $|M(G)| \leq p^{\frac{1}{2}(n+k-2)(n-k-1)}$ for $p$-groups $G$ of nilpotency class $c \geq 3$ and $p \neq 3$.*

So the natural question arises here which is the following:
**Question:** Does there exist finite $p$-groups of arbitrary nilpotency class for which this new bound is attained?
The answer of this question is yes for nilpotency class 3 and 4, see Section 3, Example1 and Example2. We conclude this section by defining some notations. We say a group $G$ of order $p^n$ is of maximal class if the nilpotency class of $G$ is $n-1$. By $ES_p(p^3)$ we denote the extra-special $p$-group of order $p^3$ having exponent $p$. By $\mathbb{Z}_p^{(k)}$ we denote $\mathbb{Z}_p \times \mathbb{Z}_p \times \cdots \times \mathbb{Z}_p (k$ times). $\gamma_i(G)$ denotes the $i$-th term of the lower central series of group $G$ and $G^{ab}$ denotes the quotient group $G/\gamma_2(G)$.

## 2. Preliminaries

In this section we recall some results which will be used to prove our main result.

**Theorem 2.1.** ([15, Theorem 4.1]) *Let $G$ be a finite group and $K$ a central subgroup of $G$. Set $A = G/K$. Then $|M(G)||G' \cap K|$ divides $|M(A)||M(K)||A^{ab} \otimes K|$.*

**Theorem 2.2.** ([6, Main Theorem]) *Let $G$ be a non-abelian finite $p$-group of order $p^n$. If $|G'| = p^k$, then we have*

$$|M(G)| \leq p^{\frac{1}{2}(n+k-2)(n-k-1)+1}.$$

*In particular,*

$$|M(G)| \leq p^{\frac{1}{2}(n-1)(n-2)+1},$$

*and the equality holds in this last bound if and only if $G = H \times Z$, where $H$ is an extra special $p$-group of order $p^3$ and exponent $p$, and $Z$ is an elementary abelian $p$-group.*

The following result gives the classification of $p$-groups $G$ of class 2 such that $|M(G)|$ attains the bound.

**Theorem 2.3.** ([16, Theorem 1.1]) *Let $G$ be a finite $p$-group of order $p^n$ and nilpotency class 2 with $|G'| = p^k$. Then $|M(G)| = p^{\frac{1}{2}(n+k-2)(n-k-1)+1}$ if and only if $G$ is one of the following groups.*

*(i)* $G_1 = ES_p(p^3) \times \mathbb{Z}_p^{(n-3)}$, *for an odd prime* $p$.

*(ii)* $G_2 = \langle \alpha, \alpha_1, \alpha_2, \beta_1, \beta_2 \mid [\alpha_i, \alpha] = \beta_i, \alpha^p = \alpha_i^p = \beta_i^p = 1 (i = 1, 2) \rangle$, *for an odd prime* $p$.

*(iii)* $G_3 = \langle \alpha_1, \beta_1, \alpha_2, \beta_2, \alpha_3, \beta_3 \mid [\alpha_1, \alpha_2] = \beta_3, [\alpha_2, \alpha_3] = \beta_1, [\alpha_3, \alpha_1] = \beta_2, \alpha_i^{(p)} = \beta_i^p = 1 (i = 1, 2, 3) \rangle$, *for an odd prime* $p$.

The following result is a consequence of [13, Main Theorem].

**Theorem 2.4.** *There is no group* $G$ *of order* $p^n$ *and of class* $c \geq 3$ *such that* $|M(G)| = p^{\frac{1}{2}n(n-1)-(n+1)}$ *where* $n \geq 6$, $p$ *is odd.*

The following result immediately follows from [9, Theorem 21].

**Theorem 2.5.** *There is no group* $G$ *of order* $p^n$ *and of class* $c \geq 3$ *such that* $|M(G)| = p^{\frac{1}{2}n(n-1)-(n-1)}$.

Let $G$ be a finite $p$-group of nilpotency class 3 with centre $Z(G)$. Set $\bar{G} = G/Z(G)$. Define a homomorphism $\psi_2 : \bar{G}^{ab} \otimes \bar{G}^{ab} \otimes \bar{G}^{ab} \to \frac{\gamma_2(G)}{\gamma_3(G)} \otimes \bar{G}^{ab}$ such that $\psi_2(\bar{x}_1 \otimes \bar{x}_2 \otimes \bar{x}_3) = [x_1, x_2]_\gamma \otimes \bar{x}_3 + [x_2, x_3]_\gamma \otimes \bar{x}_1 + [x_3, x_1]_\gamma \otimes \bar{x}_2$, where $\bar{x}$ denotes the image in $\bar{G}$ of the element $x \in G$ and $[x, y]_\gamma$ denotes the image in $\frac{\gamma_2(G)}{\gamma_3(G)}$ of the commutator $[x, y] \in G$. Define another homomorphism $\psi_3 : \bar{G}^{ab} \otimes \bar{G}^{ab} \otimes \bar{G}^{ab} \otimes \bar{G}^{ab} \to \gamma_3(G) \otimes \bar{G}^{ab}$ such that
$\psi_3(\bar{x}_1 \otimes \bar{x}_2 \otimes \bar{x}_3 \otimes \bar{x}_4) = [[x_1, x_2], x_3] \otimes \bar{x}_4 + [x_4, [x_1, x_2]] \otimes \bar{x}_3 + [[x_3, x_4], x_1] \otimes \bar{x}_2$
$+[x_2, [x_3, x_4]] \otimes \bar{x}_1$.

**Theorem 2.6.** ([11, Proposition 1] *and* [10]) *Let* $G$ *be a finite* $p$*-group of nilpotency class* 3. *With the notations above, we have*

$$|M(G)||\gamma_2(G)||Image(\psi_2)||Image(\psi_3)| \leq |M(G^{ab})||\tfrac{\gamma_2(G)}{\gamma_3(G)} \otimes \bar{G}^{ab}||\gamma_3(G) \otimes \bar{G}^{ab}|.$$

## 3. Proof of Main Theorem

In this section we prove our main theorem. Before going to the proof of main theorem, we need some reults. We use Theorem 2.2 and Theorem 2.1 in the proof of the following lemma without further reference.

**Lemma 3.1.** *Let* $G$ *be a non-abelian* $p$*-group of order* $p^n$ *with* $|G'| = p^k$ *and* $|M(G)|$ *attains the bound. Then the following hold:*

*(i)* $G^{ab}$ *is an elementary abelian* $p$*-group.*

*(ii)* $Z(G)$ *is an elementary abelian* $p$*-group.*

*(iii)* $Z(G) \subseteq G'$ (*except* $G \cong ES_p(p^3) \times \mathbb{Z}_p^{(n-3)}$).

*Proof.* *(i)* Let $K$ be a central subgroup of order $p$ such that $K \subseteq G'$. Now $|G/K| = p^{n-1}$ and $|(G/K)'| = p^{k-1}$. Now we get

$$|M(G/K)| \leq p^{\frac{1}{2}(n-1+k-1-2)(n-1-k+1-1)+1} = p^{\frac{1}{2}(n+k-4)(n-k-1)+1}.$$

Hence the inequality $|M(G)|p \leq |M(G/K)||G^{ab}|$ gives

$$|M(G)| \leq p^{\frac{1}{2}(n+k-4)(n-k-1)+1}p^{(n-k-1)} = p^{\frac{1}{2}(n+k-2)(n-k-1)+1}.$$

It follows that $|M(G/K)|$ attains the bound and $G/G'$ is elementary abelian $p$-group.

$(ii)$ Suppose the exponent of $Z(G) > p$. Consider a cyclic central subgroup $K$ of order $p^2$. Either $K \subset G'$, $K \cap G' = 1$ or $K \cap G' = \mathbb{Z}_p$.

For the first case $|G/K| = p^{n-2}$ and $|(G/K)'| = p^{k-2}$. Hence

$$|M(G/K)| \leq p^{\frac{1}{2}(n-2+k-2-2)(n-2-k+2-1)+1} = p^{\frac{1}{2}(n+k-6)(n-k-1)+1}.$$

Therefore from $|M(G)|p^2 \leq |M(G/K)||G^{ab}|$ we have

$$|M(G)| \leq p^{\frac{1}{2}(n+k-4)(n-k-1)+1}p^{(n-k-2)} = p^{\frac{1}{2}(n+k-2)(n-k-1)+1-(n-k)},$$

which is a contradiction. Other cases follows similarly.

$(iii)$ Suppose $Z(G) \nsubseteq G'$, consider a central subgroup $K$ of order $p$ such that $K \cap G' = 1$. Since $|G/K| = p^{n-1}$ and $|(G/K)'| = p^k$. So

$$|M(G/K)| \leq p^{\frac{1}{2}(n+k-3)(n-k-2)+1}.$$

Hence we have

$$\begin{aligned}|M(G)| \leq |M(G/K)||G/G'K| &\leq p^{\frac{1}{2}(n+k-3)(n-k-2)+1}p^{(n-k-1)} \\ &= p^{\frac{1}{2}(n+k-2)(n-k-1)+1-(k-1)},\end{aligned}$$

which is a contradiction for $k > 1$. For $k = 1$, $|M(G)|$ attains the bound if and only if $G \cong ES_p(p^3) \times \mathbb{Z}_p^{(n-3)}$. □

The following lemma follows from the proof of Lemma 3.1$(i)$ using Lemma 3.1$(ii)$.

**Lemma 3.2.** *If $G$ is a $p$-group of order $p^n$ such that $|M(G)|$ attains the bound, then for every central subgroup $K$ of order $p$, $|M(G/K)|$ also attains the bound.*

**Lemma 3.3.** *There is no group $G$ of order $p^n$ $(n \geq 4)$ having maximal class such that $|M(G)|$ attains the bound.*

*Proof.* First we prove that $|M(G)| \leq p^{n-2}$ for $p$-groups $G$ of maximal class. We use induction argument on $n$ to prove this.

Let $n = 4$. Then for $p = 2$ using HAP[18] of GAP[19] and for odd $p$ by [5, page. 4177] it follows that $|M(G)| \leq p^2 = p^{n-2}$.

Now consider group $G$ of order $p^n$ $(n > 4)$ of maximal class. Note that $G/Z(G)$ is also of maximal class. So by induction hypothesis $|M(G/Z(G))| \leq p^{n-3}$. Hence it follows from [14, Proposition 2.4] that

$$|M(G)|p \leq |M(G/Z(G))||G^{ab}| \leq p^{n-1}.$$

Hence $|M(G)| \leq p^{n-2}$. Now if $|M(G)|$ attains the bound for $p$-groups $G$ of maximal class then $|M(G)| = p^{\frac{1}{2}(n+n-2-2)(n-n+2-1)+1} = p^{n-1}$, which is a contradiction. □

In view of Lemma 3.2 we observe that it is sufficient to consider groups $G/K$ such that $|M(G/K)|$ attains the bound for every central subgroup $K$ of order $p$. This observation is going to be key ingredient in the proof. The following lemma refutes the existence of finite $p$-groups $G$ such that $G/K$ is of nilpotency class 2 and $|M(G)|$ attains the bound.

**Lemma 3.4.** *There is no non-abelian $p$-group $G$ of order $p^n$, $p \neq 3$ such that $G/K$ is of nilpotency class 2 for some central subgroup $K$ of order $p$ and $|M(G)|$ attains the bound. For $p = 3$, there is a group $G$ such that $|M(G)|$ attains the bound.*

*Proof.* Suppose that $G$ is a group of order $p^n$ and $|G'| = p^k$ such that $|M(G)|$ attains the bound. Let $G/K$ is of class 2 for a central subgroup $K$ of order $p$. By Lemma 3.2 $|M(G/K)|$ also attains the bound. Hence by Theorem 2.3 we have $G/K \cong G_1, G_2$ or $G_3$. Now we consider the cases depending on the structure of $G/K$.

If $G/K \cong G_1 = ES_p(p^3) \times \mathbb{Z}_p^{(n-3)}$, then $k = 2$ and $|M(G)| = p^{\frac{1}{2}(n+k-2)(n-k-1)+1}$ $= p^{\frac{1}{2}n(n-1)-(n-1)}$, which contradicts Theorem 2.5. If $G/K \cong G_2$ then $k = 3$ and $|M(G)| = p^8 = p^{\frac{1}{2}6(6-1)-(6+1)}$, which contradicts Theorem 2.4.

If $G/K \cong G_3 = \langle \alpha_1, \beta_1, \alpha_2, \beta_2, \alpha_3, \beta_3 \mid [\alpha_1, \alpha_2] = \beta_3, [\alpha_2, \alpha_3] = \beta_1, [\alpha_3, \alpha_1] = \beta_2, \alpha_i^{(p)} = \beta_i^p = 1 (i = 1, 2, 3) \rangle$, then $k = 4$ and $|M(G)| = p^{\frac{1}{2}(n+k-2)(n-k-1)+1}$ $= p^{10}$. We want to show that $|Image(\psi_3)| \geq p$ for $p \neq 3$. Then from Theorem 2.6 we have $|M(G)| \leq p^9$, which is a contradiction.

Suppose $[\beta_i, \alpha_j]$ is non-trivial in $\gamma_3(G)$ for some $j \in \{1, 2, 3\}$. Without loss of generality assume that $i = 1$. Now if $j = 2$ or 3, then $\psi_3(\alpha_2 \otimes \alpha_3 \otimes \alpha_j \otimes \alpha_1)$ is non-trivial element. Then $|Image(\psi_3)| \geq p$. Similarly we can show for $i = 2, 3$.

Hence we consider the case $[\beta_i, \alpha_i] \in \gamma_3(G)$ is non-trivial and $[\beta_i, \alpha_j] = 1$ for $i, j \in \{1, 2, 3\}, i \neq j$. Now let $|Z(G)| > p$. Without loss of generality assume that $\beta_2 = [\alpha_3, \alpha_1] \in Z(G)$ and there is a non-trivial element $[\beta_1, \alpha_1] \in \gamma_3(G)$. Then $\psi_3(\alpha_2 \otimes \alpha_3 \otimes \alpha_1 \otimes \alpha_3)$ is non-trivial element. So $|Image(\psi_3)| > p$. Next case remains $Z(G) = \langle \gamma \rangle \cong \mathbb{Z}_p$. Suppose $Image(\psi_3) = \{1\}$. Then $\psi_3(\alpha_1 \otimes \alpha_2 \otimes \alpha_3 \otimes \alpha_1) = \psi_3(\alpha_3 \otimes \alpha_1 \otimes \alpha_2 \otimes \alpha_3) = 1$ forces to have $[\beta_3, \alpha_3] = [\beta_2, \alpha_2] = [\beta_1, \alpha_1]$ and by Hall-Witt identity we have $p = 3$.

Using these relations we construct a group $G = \langle \alpha_1, \beta_1, \alpha_2, \beta_2, \alpha_3, \beta_3, \gamma \mid [\alpha_1, \alpha_2] = \beta_3, [\alpha_2, \alpha_3] = \beta_1, [\alpha_3, \alpha_1] = \beta_2, [\beta_3, \alpha_3] = [\beta_2, \alpha_2] = [\beta_1, \alpha_1] = \gamma, \alpha_i^3 = \beta_i^3 = \gamma^3 = 1 (i = 1, 2, 3) \rangle$ which is of order $3^7$. Using HAP [18] of GAP [19] we see that $|M(G)| = p^{\frac{1}{2}(n+k-2)(n-k-1)} + 1 = p^{10}$.

□

We are now ready to prove our main result.

**Proof of Theorem 1.1.** First we prove that there is no group $G$ of order $p^n$,

$p \neq 3$ of class $c = 3$ such that $|M(G)|$ attains the bound. For a central subgroup $K$ of order $p$ in $G$, $G/K$ is of class 2 or 3. If $G/K$ is of class 2, then the result follows from Lemma 3.4. Now if $G/K$ is of nilpotency class 3 for every central subgroup $K$ of order $p$, then we use induction on $n$ to prove that there is no such $G$ such that $|M(G)|$ attains the bound. For $n = 5$, $G/K$ is of maximal class, so our result is true for $n = 5$, follows from Lemma 3.3 and Lemma 3.2. Now let $G$ be a group of class 3 and $n > 5$. If there is a central subgroup of $G/K$ of order $p$ such that the factor group is of class 2, then the result follows from Lemma 3.4 and Lemma 3.2. Hence consider that for every central subgroup of $G/K$ of order $p$, the factor group of $G/K$ is of class 3 again. Since $|G/K| = p^{n-1}$, by induction hypothesis on $n$ there is no such $G/K$ such that $|M(G/K)|$ attains the bound. Hence by Lemma 3.2, result follows for class $c = 3$.

We proved our result for $c = 3$. Now we use induction argument to complete the proof for $c > 3$. Let $G$ be a $p$-group of order $p^n$ and of class $c > 3$. If nilpotency class of $G/K$ is smaller than $c$, then by induction hypothesis on $c$, there is no $G/K$ such that $|M(G/K)|$ attain the bound. Hence result follows by Lemma 3.2. Now if nilpotency class of $G/K$ is $c$ for every central subgroup $K$ of order $p$, then we use induction on $n$ to prove our result. For $n = c+2$, $G/K$ is of maximal class. So our result is true for $n = c + 2$, by Lemma 3.3 and Lemma 3.2. Let $|G| = p^n$ of class $c$ with $n > c + 2$. If there is a central subgroup of $G/K$ of order $p$ such that the factor group is of class smaller than $c$ then the result follows by induction on $c$ and by Lemma 3.2. So now for every central subgroup of $G/K$ of order $p$, the factor group of $G/K$ is of class $c$ again. As $|G/K| = p^{n-1}$, by induction hypothesis on $n$ there is no such $G/K$ such that $|M(G/K)|$ attains the bound. Hence our result follows by Lemma 3.2. This completes the proof. □

We conclude our paper by providing some examples of groups $G$ of order $p^n$, such that $|M(G)| = p^{\frac{1}{2}(n+k-2)(n-k-1)}$ where $|G'| = p^k$.

**Example1:** Consider the group $G = \langle \alpha, \alpha_1, \alpha_2, \alpha_3, \alpha_4 \mid [\alpha, \alpha_1] = \alpha_2, [\alpha_2, \alpha] = \alpha_3, [\alpha_2, \alpha_1] = \alpha_4, \alpha^p = \alpha_i^p = 1 (i = 1, 2, 3, 4)\rangle$ from [17]. This is group of order $p^5$ with $|G'| = p^3$. Nilpotency class of $G$ is 3. For $p = 5, 7, 11, 13, 17$ using HAP [18] of GAP [19] we obtain $M(G) \cong \mathbb{Z}_p \times \mathbb{Z}_p \times \mathbb{Z}_p$. Note that $|M(G)| = p^{\frac{1}{2}(5+3-2)(5-3-1)} = p^3$.

**Example2:** Consider the group $G = \langle \alpha, \alpha_1, \alpha_2, \alpha_3, \alpha_4 \mid [\alpha_i, \alpha] = \alpha_{i+1}, \alpha^p = \alpha_1^{(p)} = \alpha_{i+1}^{(p)} = 1 (i = 1, 2, 3)\rangle$ from [17] and $\alpha_{i+1}^{(p)} = \alpha_{i+1}^p \alpha_{i+2}^{\binom{p}{2}} \cdots \alpha_{i+k}^{\binom{p}{k}} ... \alpha_{i+p}$. This is group of order $p^5$ with $|G'| = p^3$. Nilpotency class of $G$ is 4. For $p = 5, 7, 11, 13, 17$ using HAP [18] of GAP [19] we obtain $M(G) \cong \mathbb{Z}_p \times \mathbb{Z}_p \times \mathbb{Z}_p$. Note that $|M(G)| = p^{\frac{1}{2}(5+3-2)(5-3-1)} = p^3$.

**Acknowledgement**: I am very grateful to my supervisor Dr. Manoj K. Yadav for his encouragement and discussions. Also I am thankful to Dr. Pradeep K. Rai for helpful discussions. I wish to thank the Harish-Chandra Research Institute, the Dept. of Atomic Energy, Govt. of India, for providing excellent research facility.

## References

[1] Schur, I, *Uber die Darstellung der endlichen Gruppen durch gebrochene lineare Substitutionen*, J. Reine Angew. Math, **127**, (1904), 20-50.
[2] Green, J.A., *On the number of automorphisms of a finite group*, Proc. Roy. Soc, A **237**, (1956), 574-581.
[3] Berkovich, Ya. G., *On the order of the commutator subgroups and the Schur multiplier of a finite p-group.*, J. Algebra, **34** (1991), 613-637.
[4] Zhou, X., *On the order of Schur multipliers of finite p-groups,* Comm. Algebra, **22**, (1994), 1-8.
[5] Ellis, G.,*On the Schur multiplier of p-groups.*, Comm. Algebra, (1999), **27**(9): 4173-4177.
[6] Niroomand, P, *On the order of Schur multiplier of non-abelian p groups*, Journal of Algebra, **322**, (2009), 4479-4482.
[7] Niroomand, P., *Characterizing finite p-groups by their Schur multipliers*, C. R. Math. Acad. Sci. Paris, Ser, **350**, (2012), 867-870.
[8] Niroomand, P., *Characterizing finite p-groups by their Schur multipliers,* $t(G) = 5$, arXiv:1001.4257.
[9] Niroomand, P., *A note on the Schur multiplier of groups of prime power order*, Ricerche di Matematica, **61** (2012), 341-346
[10] Ellis, G., *A Bound for the Derived and Frattini Subgroups of a Prime-Power Group*, Proceedings of the American Mathematical Society, **126** (1998) 2513-2523.
[11] Ellis, G., Wiegold, J., *A bound on the Schur multiplier of a prime-power group*, Bull. Austral.math.soc, **60** (1999), 191-196.
[12] Hatui Sumana, *Classification of p-groups by their Schur multiplier*, arXiv:1605.01849.
[13] Hatui Sumana, *A characterization of finite p-groups by their Schur multiplier*, arXiv:1608.06322.
[14] Jones, M.R., *Some inequalities for multiplicators of a finite group II*, Proc. Amer. Math. Soc. **45**(2) (1974), 167-172.
[15] Jones, M.R., *Some inequalities for multiplicators of a finite group*, Proc. Amer. Math. Soc, **39**(3) (1973), 450-456.
[16] Rai, Pradeep K. *On classification of groups having Schur multiplier of maximum order*, Arch. Math., (2016).
[17] James, R., *The groups of order* $p^6$(p an odd prime), Math. Comp, **34** (1980), 613-637.
[18] Ellis, G. (2008).*HAP-Homological algebra programming* A refreed GAP 4 package (GAP Group 2005), available at http://hamilton.nuigalway.ie/Hap/www.
[19] The GAP Group. (2005). *GAP-Groups Algorithms and Programming.* version 4.4. Available at: http://www.gap-system.org.

School of Mathematics, Harish-Chandra Research Institute, Chhatnag Road, Jhunsi, Allahabad 211019, INDIA

& Homi Bhabha National Institute, Training School Complex, Anushakti Nagar, Mumbai 400085, India

*E-mail address*: sumanahatui@hri.res.in, sumana.iitg@gmail.com